
anti-E2F1

Cat #: HM1122
Rabbit polyclonal IgG
0.2 µg/µ, store at 4 °C

For research use only

BACKGROUND

E2F transcription factor 1 was originally identified through its role in transcriptional activation of the adenovirus E2 promoter. It binds to the promoter regions of many genes that control cell cycle. E2F-1 is a member of a broader family of transcription regulators including E2F-2, E2F-3, E2F-4, E2F-5 and E2F-6, each of which forms heterodimers with a second protein, DP-1.

SPECIFICITY

This antibody specifically recognizes E2F1 of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunostaining.

IMMUNOGEN

A peptide mapping at the carboxy terminus of E2F1 of human origin.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCE

1. Chellappan, S., Hiebert, S., Mudryj, M., Horowitz, J.M., and Nevins, J.R. 1991. The E2F transcription factor is a cellular target for the RB protein. *Cell* 65: 1053-1061.
2. Helin, K., Lees, J.A., Vidal, M., Dyson, N., Harlow, E., and Fattaey, A. 1992. A cDNA encoding a pRB-binding protein with properties of the transcription factor E2F. *Cell* 70: 337-350.
3. Wu L, Timmers C, Maiti B, Saavedra HI, Sang L, Chong GT, Nuckolls F, Giangrande P, Wright FA, Field SJ, Greenberg ME, Orkin S, Nevins JR, Robinson ML, Leone G. (2001) The E2F1-3 transcription factors are essential for cellular proliferation. *Nature*. 414:457-62.
4. Martin K, Trouche D, Hagemeyer C, Sorensen TS, La Thangue NB, Kouzarides T. (1995) Stimulation of E2F1/DP1 transcriptional activity by MDM2 oncoprotein. *Nature*. 375:691-4.
5. Johnson DG, Schwarz JK, Cress WD, Nevins JR. (1993) Expression of transcription factor E2F1 induces quiescent cells to enter S phase. *Nature*. 365:349-52.
6. Helin, K., Wu, C., Fattaey, A.R., Lees, J.A., Dynlacht, B.D., Ngwu, C., and Harlow, E. 1993. Heterodimerization of the transcription factors E2F-1 and DP-1 leads to cooperative transactivation. *Genes and Dev.* 7: 1850-1861.
7. Krek, W., Livingston, D.M., and Shirodkar, S. 1993. Binding to DNA and the retinoblastoma gene product promoted by complex formation of different E2F family members. *Science* 262: 1557-1560.

8. Beijersbergen, R.L., Kerkhoven, R.M., Zhu, L., Carlée, L., Voorhoeve, P.M., and Bernards, R. 1994. E2F-4, a new member of the E2F gene family, has oncogenic activity and associates with p107 *in vivo*. *Genes and Dev.* 8: 2680-2690.
9. Ginsberg, D., Vairo, G., Chittenden, T., Xiao, Z.-X., Xu, G., Wydner, K.L., DeCaprio, J.A., Lawrence, J.B., and Livingston, D.M. 1994. E2F-4, a new member of the E2F transcription factor family, interacts with p107. *Genes and Dev.* 8: 2665-2679.
10. Trimarchi, J.M., Fairchild, B., Verona, R., Moberg, K., Andon, N., and Lees, J.A. 1998. E2F-6, a member of the E2F family that can behave as a transcriptional repressor. *Proc. Natl. Acad. Sci. USA* 95: 2850-2855.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com