
anti-GATA-2

Cat #: HM1162
Mouse monoclonal IgG
0.2 µg/µl, store at 4 °C

For research use only

BACKGROUND

GATA2 is a member of a family comprising GATA1, GATA2, GATA3, GATA4, GATA5, GATA6 and OSP1. They share a conserved zinc finger DNA binding domain and are capable of binding the WGATAR consensus sequence. The expression patterns of the various GATA transcription factors may overlap. It is not yet clear how the GATA factors are able to discriminate in binding their appropriate target sites. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as in erythroid cells including early erythroid cells, mast cells, and megakaryocytes. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1.

SPECIFICITY

This antibody specifically reacts with GATA-2 of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunohistochemistry.

Molecular Weight GATA-2: 50 kDa

Western blotting positive controls: Jurkat cells.

IMMUNOGEN

A synthetic peptide at the N-terminus of human GATA-2 protein.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCES

1. Ko, L.J. and Engel, J.D. 1993. DNA-binding specificities of the GATA transcription factor family. *Mol. Cell. Biol.* 13: 4011-4022.
2. Dorfman, D.M., Wilson, D.B., Bruns, G.A., and Orkin, S.H. 1992. Human transcription factor GATA-2. Evidence for regulation of preproendothelin-1 gene expression in endothelial cells. *J. Biol. Chem.* 267: 1279-1285.
3. Laverriere, A.C., MacNeill, C., Mueller, C., Poelmann, R.E., Burch, J.B., and Evans, T. 1994. GATA-4/5/6, a subfamily of three transcription factors transcribed in developing heart and gut. *J. Biol. Chem.* 269: 23177-23184.
4. Suzuki, E., Evans, T., Lowry, J., Truong, L., Bell, D.W., Testa, J.R., and Walsh, K. 1996. The human GATA-6 gene: structure, chromosomal location, and regulation of expression by tissue-specific and mitogen-responsive signals. *Genomics* 38: 283-290.
5. Zhang, P., Behre, G., Pan, J., Iwama, A., Wara-Aswapati, N., Radomska, H.S., Auron, P.E., Tenen, D.G. and Sun, Z. (1999) Negative cross-talk between hematopoietic regulators:

GATA proteins repress PU.1. *Proc. Natl. Acad. Sci. U.S.A.* 96, 8705-8710.

6. Ozawa, Y., Towatari, M., Tsuzuki, S., Hayakawa, F., Maeda, T., Miyata, Y., Tanimoto, M. and Saito, H. (2001) Histone deacetylase 3 associates with and represses the transcription factor GATA-2. *Blood* 98, 2116-2123.
7. Tsuzuki, S., Kitajima, K., Nakano, T., Glasow, A., Zelent, A. and Enver, T. (2004) Cross talk between retinoic acid signaling and transcription factor GATA-2. *Mol. Cell. Biol.* 24 (15), 6824-6836.
8. Tong, Q., Tsai, J., Tan, G., Dalgin, G. and Hotamisligil, G.S. (2005) Interaction between GATA and the C/EBP family of transcription factors is critical in GATA-mediated suppression of adipocyte differentiation. *Mol. Cell. Biol.* 25, 706-715.

PRODUCTS FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com