
anti-Histone H1

Cat #: HM1407
Mouse polyclonal IgG
0.2 µg/µl, store at 4 °C

For research use only

BACKGROUND

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. Phosphorylation of histone H1 is thought to be involved in this process, although the exact nature of this role has yet to be elucidated. Evidence suggests that histone H1 is a part of a general repressor mechanism for stable repression of transcription, but it can also activate transcription of specific genes.

SPECIFICITY

This antibody reacts Histone H1 from a broad range of mammalian species.

It can be used in Western Blotting, immunoprecipitation and immunohistochemistry (including paraffin-embedded sections).

Molecular weight of Histone H1: 32-33 kDa.

Positive controls: HeLa cells

IMMUNOGEN

Human leukemia biopsies.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCE

1. Hohmann, P. 1983. Phosphorylation of H1 histones. *Mol. Cell Biochem.* 57: 81-92.
2. Doenecke, D. and Tonjes, R. (1986) Differential distribution of lysine and arginine residues in the closely related histones H1 and H5. Analysis of a human H1 gene. *J. Mol. Biol.* 187, 461-464.
3. Doenecke, D., et al. 1988. The H1 and core histone subtypes: differential gene expression and varied primary structures. *Adv. Enzyme Regul.* 27: 107-120.
4. Nilsson, P., et al. 1992. DNA binding of histone H1 is modulated by nucleotides. *FEBS Letts.* 313: 67-70.
5. Roth, S.Y., et al. 1992. Chromatin condensation: does histone H1 dephosphorylation play a role? *Trends Biochem. Sci.* 17: 93-98.

6. Albig, W., Drabent, B., Kunz, J., Kalff-Suske, M., Grzeschik, K.H. and Doenecke, D. (1993) All known human H1 histone genes except the H1(0) gene are clustered on chromosome 6. *Genomics* 16, 649-654
7. Wolffe, A.P. 1997. Histone H1. *Int. J. Biochem. Cell Biol.* 29: 1463-1466.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com