

**IκB-β**

Cat #: HM1243  
Rabbit polyclonal IgG  
0.2 μg/μl, store at 4 °C

For research use only

**BACKGROUND**

NF-κB transcription factors are a family of structurally-related proteins that are involved in the control of a variety of cellular processes, such as growth, development, and the inflammatory response. The activity of NF-κB is tightly regulated by interaction with inhibitory IκB proteins. IκB family of proteins comprises four groups: IκB-α, IκB-β, IκB-γ, and IκB-ε. IκB proteins inactivate NFκB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IκB proteins by kinases (IκBKA, or IκBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NFκB complex. IκB-α and IκB-β mediate different NFκB responses. IκB-α appears to control more transient activation of NFκB in response to an inducer, while IκB-β controls a persistent response.

**SPECIFICITY**

This antibody reacts with IκB-β of mouse, rat and human origin by Western blotting, immunoprecipitation and immunohistochemistry.

Recommended dilution for Western blotting: 1:1000. Molecular weight of IκB-β: 45 kDa.

**IMMUNOGEN**

A peptide at the carboxy terminus of mouse IκB-β.

**STORAGE**

This antibody is stable for 12 months when stored at 2-8°C.

**REFERENCES**

1. Ghosh, S. and Baltimore, D. 1990. Activation *in vitro* to NFκB by phosphorylation of its inhibitor IκB. *Nature* 344: 678-682.
2. Kerr, L.D., Inoue, J.-I., Davis, N., Link, E., Baeuerle, P.A., Bose, H.A.J., and Verma, I.M. 1991. The Rel-associated pp40 protein prevents DNA binding of Rel and NFκB: relationship with IκB-β and regulation by phosphorylation. *Genes and Dev.* 5: 1464-1476.
3. Davis, N., Ghosh, S., Simmons, D.L., Tempst, P., Liou, H.-C., Baltimore, D., and Bose, H.R. 1991. Rel-associated pp40: an inhibitor of the Rel family of transcription factors. *Science* 252: 1268-1271.
4. Haskill, S., Beg, A.A., Tompkins, S.M., Morris, J.S., Yurochko, A.D., Sampson-Johannes, A., Mondal, K., Ralph, P., and Baldwin, A.S. 1991. Characterization of an immediate-early gene induced in adherent monocytes that encodes IκB-like activity. *Cell* 65: 1281-1289.

5. Inoue, J.-I., Kerr, L.D., Kakizuka, A., and Verma, I.M. 1992. IκB-γ, a 70 kd protein identical to the C-terminal half of p110 NFκB; a new member of the IκB family. *Cell* 68: 1109-1120.
6. Thompson, J.E., Phillips, R.J., Erdjument-Bromage, H., Tempst, P., and Ghosh, S. 1995. IκB-β regulates the persistent response in biphasic activation of NFκB. *Cell* 80: 573-582.
7. Whiteside, S.T., Epinat, J.C., Rice, N.R., and Israel, A. 1997. IκB-β, a novel member of the IκB family, controls RelA and cRel NF-κB activity. *EMBO J.* 16: 1413-1426.
8. Simeonidis, S., Liang, S., Chen, G., and Thanos, D. 1997. Cloning and functional characterization of mouse IκB-ε. *Proc. Natl. Acad. Sci. USA* 94: 14372-14377.

**PRODUCT FROM HYPROMATRIX, INC.****A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™  
Catalog Number HM3000
2. Apoptosis AntibodyArray™  
Catalog Number HM4000
3. Cell Cycle AntibodyArray™  
Catalog Number HM5000

**B. Staining AntibodyArray™s**

1. Staining AntibodyArray™ I  
Catalog Number HM8100
2. AntibodyArray Staining Apparatus  
Catalog Number HM8000

**C. Antibodies****1. HRP-conjugated antibodies**

- anti-phosphotyrosine  
Catalog Number HM2040
- anti-phosphoserine  
Catalog Number HM2070
- anti-phosphothreonine  
Catalog Number HM2090

and more...

**2. Primary antibodies**

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at [www.hypromatrix.com](http://www.hypromatrix.com).

**CONTACT**

**Hypromatrix, Inc.**  
100 Barber Avenue  
Worcester, MA 01606  
USA

Tel: 508-856-7900  
Fax: 508-302-0748  
Email: [contact@hypromatrix.com](mailto:contact@hypromatrix.com)

