

anti-Stat5a

Cat #: HM1358
Rabbit polyclonal IgG
0.2 µg/µl, store at 4 °C

For research use only

BACKGROUND

Stats (signal transducers and activators of transcription) are transcription factors activated by peptide hormones, growth factors and cytokines, particularly interferons. STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5 is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis.

SPECIFICITY

This antibody specifically recognizes Stat5a of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunostaining.

IMMUNOGEN

Recombinant protein corresponding to the Carboxyl-terminal human Stat5a protein.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCES

1. Darnell JE Jr. 1997. STATs and gene regulation. *Science* 277:1630-5.
2. Darnell, J.E., Kerr, I.M., and Stark, G.R. 1994. Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* 264: 1415-1421.
3. Pallard, C., Gouilleux, F., Charon, M., Groner, B., Gisselbrecht, S., and Dusanter-Fourt, I. 1995. Interleukin-3, erythropoietin, and prolactin activate a STAT5-like factor in lymphoid cells. *J. Biol. Chem.* 270: 15942-15945.
4. Stocklin, E., Wissler, M., Gouilleux, F. and Groner, B. (1996) Functional interactions between Stat5 and the glucocorticoid receptor. *Nature* 383, 726-728
5. Socolovsky, M., Fallon, A.E., Wang, S., Brugnara, C. and Lodish, H.F. (1999) Fetal anemia and apoptosis of red cell progenitors in Stat5a^{-/-}5b^{-/-} mice: a direct role for Stat5 in Bcl-X(L) induction. *Cell* 98, 181-191.
6. Leong, P.L., Xi, S., Drenning, S.D., Dyer, K.F., Wentzel, A.L., Lerner, E.C., Smithgall, T.E. and Grandis, J.R. (2002) Differential function of STAT5 isoforms in head and neck cancer growth control. *Oncogene* 21, 2846-2853.
7. Litterst, C.M., Kliem, S., Marilley, D. and Pfizner, E. (2003) NCoA-1/SRC-1 is an essential coactivator of STAT5 that

binds to the FDL motif in the alpha-helical region of the STAT5 transactivation Domain. *J. Biol. Chem.* 278, 45340-45351.

8. Gewinner, C., Hart, G., Zachara, N., Cole, R., Beisenherz-Huss, C. and Groner, B. (2004) The coactivator of transcription CREB-binding protein interacts preferentially with the glycosylated form of Stat5. *J. Biol. Chem.* 279, 3563-3572.
9. Lanvin, O., Gouilleux, F., Mullie, C., Maziere, C., Fuentes, V., Bissac, E., Dantin, F., Maziere, J.C., Regnier, A., Lassoued, K. and Gouilleux-Gruart, V. (2004) Interleukin-7 induces apoptosis of 697 pre-B cells expressing dominant-negative forms of STAT5: evidence for caspase-dependent and -independent mechanisms. *Oncogene* 23, 3040-3047.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com