
Anti-TDAG51

Cat #: HM1367
Mouse monoclonal IgG
0.2 µg/µl, store at 4 °C

For research use only

BACKGROUND

TDAG51 (T cell death associated gene 51) was originally described as being required for the T cell receptor-dependent induction of Fas/CD95 expression in a murine T cell hybridoma. It can restore activation-induced apoptosis in cells that have lost the ability to display Fas in response to activation. TDAG51 may play a critical role in T cell apoptosis by coupling TCR stimulation to Fas expression. TDAG51 is specifically induced by IGF-I and plays an important role in the anti-apoptotic effects of IGF-I. TDAG51 contains pleckstrin homology domains which bind polyphosphoinositides. TDAG51 gene (PHLDA1) contains (CAG)_n trinucleotide repeat which has been associated with several neurological disorders in which the repeats code for polyglutamine in the protein.

SPECIFICITY

This antibody specifically reacts with TDAG51 of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunohistochemistry.

Molecular Weight of TDAG51: 40 kDa.
Western blotting positive controls: Hep G2 cell lysate.

IMMUNOGEN

A synthetic peptide derived from N-terminus of human TDAG51 protein.

STORAGE

This antibody is stable for 24 months when stored at 2-8°C.

REFERENCES

1. Drappa, J., Brot, N., and Elkon, K. 1993. The Fas protein is expressed at high levels on CD4+CD8+ thymocytes and activated mature lymphocytes in normal mice but not in the lupus-prone strain, MRL *lpr/lpr*. Proc. Natl. Acad. Sci. USA 90: 10340-10344.
2. Park, C.G., Lee, S.Y., Kandala, G., Lee, S.Y., and Choi, Y. 1996. A novel gene product that couples TCR signaling to Fas (CD95) expression in activation-induced cell death. Immunity 4: 583-591.
3. Frank D, Mendelsohn CL, Ciccone E, Svensson K, Ohlsson R, Tycko B. (1999) A novel pleckstrin homology-related gene family defined by *Ipl/Tssc3*, TDAG51, and *Tih1*: tissue-specific expression, chromosomal location, and parental imprinting. Mamm Genome. 10:1150-9.
4. Neef R, Kuske MA, Prols E, Johnson JP. (2002) Identification of the human PHLDA1/TDAG51 gene: down-regulation in metastatic melanoma contributes to apoptosis resistance and growth deregulation. Cancer Res. 62:5920-9.

5. Toyoshima, Y., Karas, M., Yakar, S., Dupont, J., Lee, Helman, and LeRoith, D. (2004) TDAG51 mediates the effects of insulin-like growth factor I (IGF-I) on cell survival. J. Biol. Chem. 279, 25898-25904.
6. Oberg HH, Sipos B, Kalthoff H, Janssen O, Kabelitz D. (2004) Regulation of T-cell death-associated gene 51 (TDAG51) expression in human T-cells. Cell Death Differ. 11:674-84.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com