
anti-TGF β Receptor 2

Cat #: HM1369
Rabbit polyclonal IgG
0.2 $\mu\text{g}/\mu\text{l}$, store at 4 °C

For research use only

BACKGROUND

Three members of the TGF β family, TGF β 1, TGF β 2 and TGF β 3, have been identified in mammals. TGF β s mediate their activity by high affinity binding to the type II receptor 70 kDa transmembrane protein with a cytoplasmic serine-threonine kinase domain. For signaling growth inhibition and early gene responses the type II receptor requires both its kinase activity and association with members of type I receptors called activin receptor-like kinase (ALK) 1 to 6. Of these, only ALK5 has been shown to represent a functional TGF beta type I receptor. TGF β Receptor 2 is involved in regulating cell proliferation and differentiation and extracellular matrix production and acts as a signal transducer. It phosphorylates TGFBR1 upon binding to its ligand, TGF-beta. TGFBR1 in turn phosphorylates the cytoplasmic effectors of the pathway, known as SMADs.

SPECIFICITY

This antibody reacts TGF β RII p70 of mouse, rat and human origin by Western blotting, immunoprecipitation and immunohistochemistry (including paraffin-embedded sections); non cross-reactive with TGF β RI p55.

Molecular Weight of TGF β RII: 70 kDa.

IMMUNOGEN

A peptide at an internal domain of the human TGF β RII.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCES

1. Cheifetz, S., Hernandez, H., Laiho, M., ten Dijke, P., Iwata, K.K., and Massagué, J. 1990. Distinct transforming growth factor- β receptor subsets as determinants of cellular responsiveness to three TGF- β isoforms. *J. Biol. Chem.* 265: 20533-20538.
2. Wrana, J.L., Attisano, L., Cárcamo, J., Zentella, A., Doody, J., Laiho, M., Wang, X., and Massagué, J. 1992. TGF β signals through a heteromeric protein kinase receptor complex. *Cell* 71: 1003-1014.
3. Massagué, J. 1992. Receptors for the TGF- β family. *Cell* 69: 1067-1070.
4. Attisano, L., Cárcamo, J., Ventura, F., Weis, F.M.B., Massagué, J., and Wrana, J.L. 1993. Identification of human activin and TGF β type I receptors that form heteromeric kinase complexes with type II receptors. *Cell* 75: 671-680.
5. Franzen, P., ten Dijke, P., Ichijo, H., Yamashita, H., Schulz, P., Heldin, C.H. and Miyazono, K. (1993) Cloning of a TGF beta type I receptor that forms a heteromeric complex with the TGF beta type II receptor. *Cell* 75, 681-692.
6. Franzen, P., ten Dijke, P., Ichijo, H., Yamashita, H., Schulz, P., Heldin, C., and Miyazono, K. 1993. Cloning of a TGF β

type I receptor that forms a heteromeric complex with the TGF β type II receptor. *Cell* 75: 681-692.

7. Perlman, R., Schiemann, W.P., Brooks, M.W., Lodish, H.F. and Weinberg, R.A. (2001) TGF-beta-induced apoptosis is mediated by the adapter protein Daxx that facilitates JNK activation. *Nat. Cell Biol.* 3, 708-714.
8. Loeys, B.L., Chen, J., Neptune, E.R., Judge, D.P., Podowski, M., Holm, T., Meyers, J., Leitch, C.C., Katsanis, N., Sharifi, N., Xu, F.L., Myers, L.A., Spevak, P.J., Cameron, D.E., De Backer, J., Hellemans, J., Chen, Y., Davis, E.C., Webb, C.L., Kress, W., Coucke, P., Rifkin, D.B., De Paepe, A.M. and Dietz, H.C. (2005) A syndrome of altered cardiovascular, craniofacial, neurocognitive and skeletal development caused by mutations in TGFBR1 or TGFBR2. *Nat. Genet.* 37, 275-281.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com