
Anti-TRAF5

Cat #: HM1381
Rabbit polyclonal IgG
0.2 µg/µl, store at 4 °C

For research use only

BACKGROUND

Members of the TNF receptor associated factor (TRAF) protein family have been implicated in the signal transduction mechanism of the TNF receptor superfamily. Six members have been described and are designated TRAF1, TRAF2, TRAF3, TRAF4, TRAF5, and TRAF6. TRAF5 is one of the components of a multiple protein complex associated with the CD40 cytoplasmic domain, which mediates TNF induced NF-kappa B activation and protection from cell death. TRAF5 has also been implicated in the signaling events mediated by various other receptors including CD27, CD30, and lymphotoxin-beta receptor.

SPECIFICITY

This antibody reacts with TRAF5 of mouse, rat and human origin by Western blotting, immunoprecipitation and immunohistochemistry.

Recommended dilution for Western blotting: 1:1000.

IMMUNOGEN

Full length human TRAF5 recombinant protein.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCES

1. Baker, S.J. and Reddy, E.P. 1996. Transducers of life and death: TNF receptor superfamily and associated proteins. *Oncogene* 12: 1-9.
2. Rothe, M., Sarma, V., Dixit, V.M., and Goeddel, D.V. 1995. TRAF2-mediated activation of NF-κB by TNF receptor 2 and CD40. *Science* 269: 1424-1427.
3. Cheng, G. and Baltimore, D. 1996. TANK, a co-inducer with TRAF2 of TNF- and CD 40L-mediated NF-κB activation. *Genes and Dev.* 10: 963-973.
4. Regnier, C.H., Tomasetto, C., Moog-Lutz, C., Chenard, M.P., Wendling, C., Basset, P., and Rio, M.C. 1995. Presence of a new conserved domain in CART1, a novel member of the tumor necrosis factor receptor-associated protein family, which is expressed in breast carcinoma. *J. Biol. Chem.* 270: 25715-25721.
5. Nakano, H., Oshima, H., Chung, W., Williams-Abbott, L., Ware, C.F., Yagita, H., and Okumura, K. 1996. TRAF5, an activator of NF-κB and putative signal transducer for the lymphotoxin receptor. *J. Biol. Chem.* 271: 14661-14664.
6. Ishida, T.K., Tojo, T., Aoki, T., Kobayashi, N., Ohishi, T., Watanabe, T., Yamamoto, T. and Inoue, J. (1996) TRAF5, a novel tumor necrosis factor receptor-associated factor family protein, mediates CD40 signaling. *Proc. Natl. Acad. Sci. U.S.A.* 93, 9437-9442.

7. Nakano, H., Sakon, S., Koseki, H., Takemori, T., Tada, K., Matsumoto, M., Munechika, E., Sakai, T., Shirasawa, T., Akiba, H., Kobata, T., Santee, S.M., Ware, C.F., Rennert, P.D., Taniguchi, M., Yagita, H. and Okumura, K. (1999) Targeted disruption of Traf5 gene causes defects in CD40- and CD27-mediated lymphocyte activation. *Proc. Natl. Acad. Sci. U.S.A.* 96, 9803-9808.
8. Tada, K., Okazaki, T., Sakon, S., Kobarai, T., Kurosawa, K., Yamaoka, S., Hashimoto, H., Mak, T.W., Yagita, H., Okumura, K., Yeh, W.C. and Nakano, H. (2001) Critical roles of TRAF2 and TRAF5 in tumor necrosis factor-induced NF-kappa B activation and protection from cell death. *J. Biol. Chem.* 276, 36530-36534.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com