
Anti-Tuberin

Cat #: HM1387
Rabbit polyclonal IgG
0.2 µg/µl, store at 4 °C

For research use only

BACKGROUND

Tuberous sclerosis (TSC) is a genetic disorder characterized by mental retardation and widespread benign and infrequently malignant tumors in a variety of tissues. Two different genetic loci have been linked to TSC; TSC2 encodes a 180 kDa protein, 1784 amino acids in length, called tuberin. Tuberin exhibits a region of limited homology to the catalytic domain of Rap1 GAP. It is capable of stimulating the intrinsic GTPase activity of Rap 1A. Tuberin is believed to be a tumor suppressor. It associates with hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin.

SPECIFICITY

This antibody specifically reacts with Tuberin of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunohistochemistry.

IMMUNOGEN

A synthetic peptide derived from N-terminus of human Tuberin protein.

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCES

1. The European Chromosome 16 Tuberous Sclerosis Consortium. 1993. Identification and characterization of the tuberous sclerosis gene on chromosome 16. *Cell* 75: 1305-1315.
2. Wienecke, R., Konig, A., and DeClue, J.E. 1995. Identification of tuberin, the tuberous sclerosis-2 product. Tuberin possesses specific Rap1GAP activity. *J. Biol. Chem.* 270: 16409-16414.
3. Carsillo, T., Astrinidis, A. and Henske, E.P. (2000) Mutations in the tuberous sclerosis complex gene TSC2 are a cause of sporadic pulmonary lymphangiomyomatosis. *Proc. Natl. Acad. Sci. U.S.A.* 97, 6085-6090.
4. Inoki, K., Li, Y., Zhu, T., Wu, J. and Guan, K.L. (2002) TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR Signaling. *Nat. Cell Biol.* 4, 648-657.
5. Shumway, S.D., Li, Y. and Xiong, Y. (2003) 14-3-3beta binds to and negatively regulates the tuberous sclerosis complex 2 (TSC2) tumor suppressor gene product, tuberin. *J. Biol. Chem.* 278 (4), 2089-2092.
6. El-Hashemite, N., Zhang, H., Henske, E.P. and Kwiatkowski, D.J. (2003) Mutation in TSC2 and activation of mammalian target of rapamycin signalling pathway in renal angiomyolipoma. *Lancet* 361, 1348-1349.

7. Astrinidis, A., Senapedis, W., Coleman, T.R. and Henske, E.P. (2003) Cell cycle-regulated phosphorylation of hamartin, the product of the tuberous sclerosis complex 1 gene, by cyclin-dependent kinase 1/cyclin B. *J. Biol. Chem.* 278, 51372-51379.
8. Li, Y., Inoki, K. and Guan, K.L. (2004) Biochemical and functional characterizations of small GTPase Rheb and TSC2 GAP activity. *Mol. Cell. Biol.* 24, 7965-7975.
9. Rosner, M. and Hengstschlager, M. (2004) Tuberin binds p27 and negatively regulates its interaction with the SCF component Skp2. *J. Biol. Chem.* 279, 48707-48715.

PRODUCT FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™
Catalog Number HM3000
2. Apoptosis AntibodyArray™
Catalog Number HM4000
3. Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

1. Staining AntibodyArray™ I
Catalog Number HM8100
2. AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT

Hypromatrix, Inc.
100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900
Fax: 508-302-0748
Email: contact@hypromatrix.com
Web: www.hypromatrix.com