

---

**anti-c-Cbl**

Cat #: HM1070  
Mouse monoclonal IgG  
0.2 µg/µl, store at 4 °C

For research use only

**BACKGROUND**

c-Cbl was identified as the cellular homolog of the murine Cas NS-1 leukemia retroviral oncogene *v-cbl*, which induces pre-B and B-myeloid leukemias. *v-Cbl* represents a severely truncated form of the cellular homolog containing only the N-terminal amino acid of c-Cbl. c-Cbl is a 120 kDa protein, located in the cytoplasm and cytoskeleton, and is widely expressed in hematopoietic cell lines. c-Cbl comprises an N-terminal transforming region (Cbl-N), which contains a phosphotyrosine binding domain, a RING zinc-finger domain adjacent to Cbl-N, and a large C-terminal region (Cbl-C) containing a large proline-rich region and a leucine zipper. The proline-rich region contains multiple potential tyrosine phosphorylation sites and docking sites for SH2 and SH3-containing proteins. c-Cbl is involved in tyrosine kinase-dependent signaling pathways. It is one of the earliest targets of tyrosine phosphorylation in response to a number of cellular stimuli including T- and B-cell receptor activation. In addition, stimulation of growth factor receptors by EGF, PDGF, NGF and FGF, results in c-Cbl tyrosine phosphorylation, in lymphoid and other cell lines. Upon receptor activation, c-Cbl interacts with SH2 and SH3 domains of several cytoplasmic signaling proteins, including the tyrosine kinases Src, Fyn, Lck, Syk, and ZAP-70, the adaptor proteins Grb2, Nck, Shc, Crk-II and CrkL. In addition, c-Cbl can specifically associate with the p85β subunit of PI3-kinase.

**SPECIFICITY**

This antibody specifically recognizes c-Cbl of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunostaining.

**IMMUNOGEN**

A peptide corresponding to the C terminus of human c-Cbl p120.

**STORAGE**

This antibody is stable for 12 months when stored at 2-8°C.

**REFERENCES**

1. Langdon, W.Y., et al. 1989. *v-cbl*, an oncogene from a dual-recombinant murine retrovirus that induces early B-lineage lymphomas. Proc. Natl. Acad. Sci. USA 86: 1168-1172.
2. Langdon, W.Y., et al. 1989. The *c-cbl* proto-oncogene is preferentially expressed in thymus and testis tissue and encodes a nuclear protein. J. Virol. 63: 5420-5424.
3. Blake, T.J., et al. 1991. The sequences of the human and mouse *c-cbl* proto-oncogenes show *v-cbl* was generated by a large truncation encompassing a proline-rich domain and a leucine zipper-like motif. Oncogene 6: 653-657.
4. Fukazawa, T., Miyake, S., Band, V. and Band, H. (1996) Tyrosine phosphorylation of Cbl upon epidermal growth factor (EGF) stimulation and its association with EGF receptor and downstream signaling proteins. J. Biol. Chem. 271, 14554-14559.

5. Smit, L. and Borst, J. (1997) The Cbl family of signal transduction molecules. Cytogenet. Cell Genet. 8, 359-379.
6. Joazeiro, C.A., Wing, S.S., Huang, H., Leverson, J.D., Hunter, T. and Liu, Y.C. (1999) The tyrosine kinase negative regulator c-Cbl as a RING-type, E2-dependent ubiquitin-protein ligase. Science 286, 309-312.
7. Davies, G.C., Ettenberg, S.A., Coats, A.O., Mussante, M., Ravichandran, S., Collins, J., Nau, M.M. and Lipkowitz, S. (2004) Cbl-b interacts with ubiquitinated proteins; differential functions of the UBA domains of c-Cbl and Cbl-b. Oncogene 23, 7104-7115.
8. Jacob, C., Cottrell, G.S., Gehringer, D., Schmidlin, F., Grady, E.F. and Bunnett, N.W. (2005) c-Cbl mediates ubiquitination, degradation, and down-regulation of human protease-activated receptor 2. J. Biol. Chem. 280, 16076-16087.

**PRODUCTS FROM HYPROMATRIX, INC.****A. AntibodyArray™s:**

1. Signal Transduction AntibodyArray™  
Catalog Number HM3000
2. Apoptosis AntibodyArray™  
Catalog Number HM4000
3. Cell Cycle AntibodyArray™  
Catalog Number HM5000

**B. Staining AntibodyArray™s**

1. Staining AntibodyArray™ I  
Catalog Number HM8100
2. AntibodyArray Staining Apparatus  
Catalog Number HM8000

**C. Antibodies****1. HRP-conjugated antibodies**

- anti-phosphotyrosine  
Catalog Number HM2040
- anti-phosphoserine  
Catalog Number HM2070
- anti-phosphothreonine  
Catalog Number HM2090

and more...

**2. Primary antibodies**

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at [www.hypromatrix.com](http://www.hypromatrix.com).

**CONTACT**

**Hypromatrix, Inc.**  
100 Barber Avenue  
Worcester, MA 01606  
USA

Tel: 508-856-7900  
Fax: 508-302-0748  
Email: [contact@hypromatrix.com](mailto:contact@hypromatrix.com)  
Web: [www.hypromatrix.com](http://www.hypromatrix.com)