

anti-p-IκB-α

Cat #: HM1242
Mouse monoclonal IgG
0.2 μg/μl, store at 4 °C

For research use only

BACKGROUND

NF-κB transcription factors are a family of structurally-related proteins that are involved in the control of a variety of cellular processes, such as growth, development, and the inflammatory response. The activity of NF-κB is tightly regulated by interaction with inhibitory IκB proteins. IκB family of proteins comprises four groups: IκB-α, IκB-β, IκB-γ, and IκB-ε. IκB proteins inactivate NFκB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IκB proteins by kinases (IκBKA, or IκBK) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NFκB complex.

SPECIFICITY

This antibody specifically recognizes Ser-32 phosphorylated IκB-α of human, mouse and rat origin.

The antibody can be used in Western blotting, immunoprecipitation and immunostaining (including paraffin-embedded sections).

Molecular weight of p- IκB-α: 37 kDa.

Western blotting positive control: TNF α -treated HeLa cells.

IMMUNOGEN

A peptide containing phosphorylated Ser-32 of IκB-α of human origin

STORAGE

This antibody is stable for 12 months when stored at 2-8°C.

REFERENCES

- Inoue, J.-I., Kerr, L.D., Kakizuka, A., and Verma, I.M. 1992. IκB-α, a 70 kd protein identical to the C-terminal half of p110 NFκB; a new member of the IκB family. *Cell* 68: 1109-1120.
- Whiteside, S.T., Epinat, J.C., Rice, N.R., and Israel, A. 1997. IκB-β, a novel member of the IκB family, controls RelA and cRel NF-κB activity. *EMBO J.* 16: 1413-1426.
- Kerr, L.D., Inoue, J.-I., Davis, N., Link, E., Baeuerle, P.A., Bose, H.A.J., and Verma, I.M. 1991. The Rel-associated pp40 protein prevents DNA binding of Rel and NFκB: relationship with IκB-α and regulation by phosphorylation. *Genes and Dev.* 5: 1464-1476.
- Davis, N., Ghosh, S., Simmons, D.L., Tempst, P., Liou, H.-C., Baltimore, D., and Bose, H.R. 1991. Rel-associated pp40: an inhibitor of the Rel family of transcription factors. *Science* 252: 1268-1271.
- Haskill, S., Beg, A.A., Tompkins, S.M., Morris, J.S., Yurochko, A.D., Sampson-Johannes, A., Mondal, K., Ralph, P., and Baldwin, A.S. 1991. Characterization of an immediate-early gene induced in adherent monocytes that encodes IκB-like activity. *Cell* 65: 1281-1289.
- Ghosh, S. and Baltimore, D. 1990. Activation *in vitro* to NF κB by phosphorylation of its inhibitor IκB. *Nature* 344: 678-682.
- Thompson, J.E., Phillips, R.J., Erdjument-Bromage, H., Tempst, P., and Ghosh, S. 1995. IκB-β regulates the persistent response in biphasic activation of NFκB. *Cell* 80: 573-582.
- Simeonidis, S., Liang, S., Chen, G., and Thanos, D. 1997. Cloning and functional characterization of mouse IκB. *Proc. Natl. Acad. Sci. USA* 94: 14372-14377.
- Prajapati, S., R. B. Gaynor. 2002. Regulation of I B kinase (IKK) /NEMO function by IKK -mediated phosphorylation. *J. Biol. Chem.* 277:24331.

PRODUCTS FROM HYPROMATRIX, INC.**A. AntibodyArray™s:**

- Signal Transduction AntibodyArray™
Catalog Number HM3000
- Apoptosis AntibodyArray™
Catalog Number HM4000
- Cell Cycle AntibodyArray™
Catalog Number HM5000

B. Staining AntibodyArray™s

- Staining AntibodyArray™ I
Catalog Number HM8100
- AntibodyArray Staining Apparatus
Catalog Number HM8000

C. Antibodies**1. HRP-conjugated antibodies**

- anti-phosphotyrosine
Catalog Number HM2040
- anti-phosphoserine
Catalog Number HM2070
- anti-phosphothreonine
Catalog Number HM2090

and more...

2. Primary antibodies

Hypromatrix offers a variety of high quality antibodies. For a complete list of antibodies and their specificities, please visit our web site at www.hypromatrix.com.

CONTACT**Hypromatrix, Inc.**

100 Barber Avenue
Worcester, MA 01606
USA

Tel: 508-856-7900

Fax: 508-302-0748

Email: contact@hypromatrix.com

Web: www.hypromatrix.com